

Institute of Meteorology and Water Management National Research Institute

DATA ASSIMILATION activities at IMWM Kraków


M. Szczęch-Gajewska, B. Bochenek





Model characteristics

	ALARO-1	
horizontal resolution	4.0km	
number of grid points	789 x 789	
vertical levels	60	
operational time step	163.6s / 120s	
coupling frequency	3h (ARPEGE LBC)	
forecast length	66h (60h at 18UTC)	
model version	CY40 T1	
HPC	Clusterof HP BL460c_GEN8 servers connected with Infiniband network, OS Scientific Linux 6, Intel Xeon E5-2690 processors – with maximum 1552 cores (97 nodes with 16 cores each), each core	
	RAM 128 GB, disc array – 64 TB.	



PL




AROME/IMWM DA

Data

for now just OPLACE base

Surface


- Data assimilation using CANARI
- . Standard setting, no special tuning
- . Test once per day run at 00UTC
 - (background our 6h forecast from 18UTC, SST copied from ARPEGE analysis, CANARI and then 66h forecast)





ALARO-1/IMWM DA - verification

HARP verification





Plans: local data

- Add local data to our CANARI run (obsoul_merge we have but not yet used)
- Start 4 times per day parallel run with CANARI
- 3D-Var ...





Plans: 3D-Var preparation

- First for AROME B matrix was built on the base of our domain AROME ensemble forecast with LBC from AEARP. AROME domain: 799x799grid points, 2.0km horizontal resolution, 60 vertical levels. Ensemble: 6 members, 2 times per day and 30 days of forecast.
- Not finished yet the same wok for ALARO-1.
 Details ask Bogdan.

