




SHMU - NWP

DA : M. Nestiak, M. Derkova, V. Tarjani, M. Imrisek, K. Catlosova *(DT)*
J. Vivoda, M. Bellus, M. Dian, O. Spaniel, R. Zehnal





Operational Setup SHMU

- ❑ on SHMU: HPC2 (RH Linux)
- ❑ ALARO-1, cy40t1_bf06
- ❑ domain: 4.5km, 625x576
- ❑ 63 vertical levels, mean orography
- ❑ time step 180s
- ❑ 3h space consistency coupling
- ❑ ARPEGE synchronous
- ❑ forecasts up to
 - ❑ F+72h at 00, 06, 12 UTC
 - ❑ F+60h at 18 UTC
- ❑ upper air blending
- ❑ CANARI (local + OPLACE)




Still running also **old** operational Setup SHMU

- ❑ on SHMU: HPC1 (AIX)
- ❑ ALARO-0, 36_t1.10
- ❑ domain: 9km, 320x288
- ❑ 37 vertical levels, mean orography
- ❑ time step 400s
- ❑ 3h space consistency coupling
- ❑ ARPEGE synchronous
- ❑ forecasts up to
 - ❑ F+72h at 00, 06, 12 UTC
 - ❑ F+60h at 18 UTC
- ❑ upper air blending
- ❑ CANARI (local + OPLACE




Tests domains (only 00, 12 UTC runs)

model: ALADIN_2km precipitation base: 2018-09-15_00 (Saturday) range: +03 valid: 2018-09-15_03 (Saturday) previous 3 hour(s)



- ALARO-1 cy40t1_bf06
- 2km, 512x384
- 73 vertical levels
- time step 90s
- Coupling with 4.5km
- HPC1-Gentoo linux - Class:old


model: AROME_2km precipitation base: 2018-09-15_00 (Saturday) range: +03 valid: 2018-09-15_03 (Saturday) previous 3 hour(s)




- AROME cy40t1_bf06
- 2km, 512x384
- 73 vertical levels
- time step 144s
- Coupling with 4.5km
- HPC2:Gentoo linux - Class:old
- B-Matrix

B-matrix: spin-up ensemble based B matrix, LBC data with 8km resolution from AEARP, 6 members, 2 periods in 2016 (32 days)

average unbal T,lnps cors



average unbal T,lnps cors



testing domain:

- AROME cy40t1_bf05
- 2km, 512x384
- 73 vertical levels

OPER domain:

- ALARO-1 cy40t1_bf05
- 4.5km, 625x576
- 63 vertical levels



Data assimilation activities with focus on GNSS data

Martin Imrišek

Slovak University of Technology in Bratislava
Department of Theoretical Geodesy


&

Slovak Hydrometeorological Institute



martin.imrisek@stuba.sk

testing domain:

- AROME cy40t1_bf05
- 2km, 512x384
- 73 vertical levels



RADAR (ODIM HDF5) -> RH.py (Slovenia) + Bator cy43t1 -> cy40t1



Testing Hobby Urban/Mountain station + NATIONAL OPLACE on ALARO-1

2km


testing domain:

- ❑ ALARO-1 cy40t1_bf06
- ❑ 2km, 512x384
- ❑ 73 vertical levels
- ❑ MESCOAN



Conclusions:

- urban stations most of time degrading MESCOAN analysis when assimilate as SYNOP (T2m, RH)
- require investigation of create new class of model observations with parameterisation of position, distance from a building, surrounding trees, height, shadow interval, ...
- very low representativeness of the surroundings in the city
- implementation in SURFEX (TEB - roof, wall, road)
 - Roof surface temperature (indoor/outdoor)
 - Wall surface temperature (indoor/outdoor)
 - ...



New SODA-EKF based assimilation suite (ViTa)

- Analysis of soil water content (WG), temperature (TG), and later also snow cover
- **Spatial domain:** INCA-SK 501 x 301 @ 1 km
- **Gridded observations:** Standard CANARI analysis is replaced by hi-res analysis of T2M & RH2M from INCA-SK system (see figure)
- **Forcing:** ~20m above surface,
 - surface fields: INCA-SK precipitation analysis + global radiation analysis -> improved calculation of Jacobians
 - other fields: from ALARO-SK 4.5km
- SURFEX and SODA executables from cy40t1 pack
- Future plans:
 - Test and optimize current setup
 - Compare EKF with OI_MAIN
 - Switch to SURFEX v8.1
 - Snow cover analysis





Fig: Downscaled CANARI (top) vs INCA-SK (bottom) analysis of T2M and RH2M used in SODA-EKF

1-column EKF validation (ViTa)

- Sensitivity test with ISBA soil scheme
- Ugly output from OFFLINE when executed as multi-process job , single-process job worked well

SURFEX v8.1 compilation (ViTa)

- SURFEX v8.1 (Master + SODA_V8 branches) downloaded from SURFEX git repository
- Issues with compilation on HPC (IBM Power7) with GNU compiler
- Need further investigation or try newer version of GNU compiler or native IBM compiler




```
/src/dir_obj-LXgfortran-SFX-V8-1-1-MPIAUTO-OMP-O2-X0/MASTER/spl1_fadcpl_fort.f90:162:0:
```

```
error: unable to generate reloads for: IF (LHOOK) CALL  
DR_HOOK('FADCPL_MT',1,ZHOOK_HANDLE)
```

```
internal compiler error: in find_reloads, at reload.c:3883
```

First results of ongoing diploma thesis on MODE -S

2 months of EHS data sample provided by the Air Traffic Control from Bratislava airport is being processed and technical tests are to start on AROME/SHMU domain asap.







Difference between Guess and Analysis at level 40. Left) Temperature Right) Wind

testing domain:

- ☐ AROME cy40t1_bf05
- ☐ 2km, 512x384
- ☐ 73 vertical levels

Future plans

- ❑ 1h RUC AROME 2km 3DVar analysis [F+12h] (GNSS, radars, HRW, automatic stations)
- ❑ 1h RUC ALARO-1 2km 3DVar analysis [F+12h](GNSS, radars, HRW, automatic stations)
- ❑ Continue works on radar DA for RC-LACE radars (Michal Nestiak)
- ❑ Continue works on GNSS data (Martin Imrisek)
- ❑ Continue works on Surfex (Viktor Tarjani)
- ❑ Calculating B-Matrix for ALARO-1 2km (Maria Derkova)
- ❑ working on MODE-S diploma thesis (Katarina Catlosova co: Maria Derkova)
- ❑ Working on Urban/Mountain automatic stations (M. Nestiak, M. Dian, R. Zehnal)



thanks for attention