

Status data assimilation in Austria


Florian Meier, Phillip Scheffknecht, Clemens Wastl, Florian Weidle, Christoph Wittmann, Stefan Schneider, Jasmin Vural




Significant changes in operations in the second sec

- C-LAEF (AROME-EPS) becomes operational end 2019
- Hourly AROME-RUC 1.2km becomes operational end 2019
- ALARO-0 4.8km will be probably switched off
- ALADIN LAEF is not full

AROME-RUC Domain & Topography





Operational configurations CMCs

ALARO-0 4.8km L60 cy40t1	AROME-Aut L90 cy40t1	C-LAEF-2.5km L90/16 cy40t1	AROME-RUC 1.2km L90 cy40t1
3h-IFS 4x/day +72h dt=180s	1h-IFS 8x/day +60h dt=60s	6h-IFS-EPS 2x/day +48h+2x6h dt=60s	1h-AROME-Aut 24x/day +12h dt=30s
dynamical downscaling	3D-VAR	3D-Var-EDA- Ens-Jk	3D-VAR +LHN+FDDA- nudging+VARQC
CANARI	CANARI-OIMAIN +MESCAN inline+ SNOW exchange/ SNOWGRID+SAT	CANARI-EDA (with screening) MESCAN +snow as AROME-Aut	CANARI-OIMAIN MESCAN +additional Tlakes +SNOW as AROME- Aut
DFI	-	-	IAU (45+5min)
	Static Ens-B from LAEF downscaling	Static Ensemble B from C-LAEF and V from IFS-EPS	Static-Ens-B from AROME-RUC EDA+ differences of the day
• •			

Observations used in AROME/C-LAEF(no sat):

Obstype	Parameter
Synop+Tawes+Ship	U10m,V10m, RH2m,T2m, Z
AMDAR	U, V, T
GEOWIND	U, V (WVCL1/2,WVMW1, IR3, VIS3)
TEMP	U, V, T, Z, Q
PILOT	U, V
MSG-SEVIRI	WV radiances
NOAA18/19/MetOp-A,-B	AMSU-A, AMSU-B, MHS, HIRS
MetOp-A	IASI
MetOp-A	U10m, V10m ASCAT ocean winds 25km

T_LAKE from Lake Constance from measurement interpolated inside OIMAIN

CANARI settings: REF_A=190km, LVARSIGO=F, LMESCAN=T, LCORRF=T REF_S_T2=5.0,REF_S_H2=0.3,RCLIMCA=0.045,RCT2SY=3.9, RCH2SY=2.5 OROLIM=3800.,ORODIF=1650. -> no change since last year




Additional observations in AROME-RUC:


Obstype	Parameter	
SYNOP national (OPLACE)	U10m,V10m, RH2m,T2m, Z	
MODE-S ARSO/CHMI (MRAR) KNMI/AT	U, V, T+AMDAR-Q	
RADAR AT/SI/D HR/CZ/SK/F (1 station)	DOW+REF +saturation of profiles	
GNSS (AT national)	ZTD	
(ATMS SUOMI+NOAA20)	Radiances ch1-3,5-14,16-22	
INCA	RR5min via LHN	
TAWES+10/20/30min	T2m/RH2m/U10m/V10m via FDDA Nudging	
T-Lake (Fertoe/Balaton) pseudo obs	TS_WATER	

CANARI settings: REF_A=110km, LVARSIGO=F, LMESCAN=T, LCORRF=T REF_S_T2=5.0,REF_S_H2=0.3,RCLIMCA=0.045,RCT2SY=3.9, -RCH2SY=2.5 OROLIM=3800.,ORODIF=1650.

AROME-Aut structure (3 hourly cycle)




AROME-RUC structure part A





Incremental Analysis Update IAU

- Idea: Bloom et al. Mon Wea Rev.1996
- "Nudging of analysis"
- adds during integration in a defined time window each timestep a part of the analysis increment to the first guess
- Implemented by MF P. Brousseau in AROME cy43t2; (backphased to cycle 40t1)
- Can be used to 1. add increment/use observations later during the forecast (MF) ->short window
- 2. or to smooth increment in time and so filter spin-up (ZAMG) ->long window
- Needs guess+analysis;
- switches: LIAU to switch IAU on, TSTARTIAU starttime of window in s ->TSTOPIAU endtime of window, ALPHAIAU amplification factor multiplied with the increment per timestep analysis










Regional Cooperation for






AROME-RUC structure part B



two hourly cycles one using minimization via IAU at 0-5min the other at 15-60min IAU with long window seems to affect hydrometeors

IAU detoriates hydrometeors Regional Cooperation for ->additional blending





S085RAIN after +15min-60min IAU S085RAIN after +1h run with LHN+FDDA-Nudg

Observation was rain=0 on that day






STD assimilation in ARQME Reference of the Control Europe (F. Weidle)

 $^{\mathsf{F}}$

nwp central europe

Operator from KNMI backphased to cy40t1 export






ROMANIA

ANM

ZAMG

GPS-RO OSSE (Phillip Scheffknecht)




Positive impact on wind and temperature above 500hPa; below mixed


windfarms in AROME-RUC (about 200 turbines)



- QC based on deviation of measured wind and power from power curve
 -> turbine in regular mode
- Blacklisting of wind turbines with large bias or standard deviation (12 turbines)
- Parameterisation of turbine effects on the environment (approach of Fitch et al. 2012)
 ->U, V, TKE tendencies based on turbine type, height, diameter and wind
- New pre-processing (Python) to handle large number of turbines
- Assimilate SCADA wind (temperature also possible, but measurement only 0 digit precision)




effect of windfarm wind assimilation on temperature




or V and T

Regional Cooperation for difference in V 6.9.2019 11 UTC+IM





reference-(assim+parameterized farms) difference due to assimilation and Parametrisation of windfarms



param-(assim+param) difference only due to assimilation




Lake temperatures from SYNOP T2m

Fertoe: Podersdorf 11084 assume 8h mean T2m=T-Lake

Regional Cooperation for

Limited Area Modeling in Central Europe

Balaton: Siofok 12935 assume 12h mean T2m=T-Lake




T2mfertoe=24°C Twaterobs=23°C T2m Balaton=24°C Twaterobs=26°C





nwp central europe


Δ CF Regional Cooperation for Limited Area Modeling in Central Europe nwp central europe


B-Matrix properties













DHM



ANM



B-Matrix properties



18



















Blacklisting aus Mittelung bzw. Quadrat der FG-Abweichungen von August 2019



"Verdächtige" Anlagen: Wind: 17820,17817, 80019, 80015, 260051 Temperatur: 701700, 701187,260071, 17818, 17819, 101148, 17812

