Regional Cooperation for Limited Area Modeling in Central Europe




#### VarBC spin up and assimilation of Slovenian GNSS ZTD observations

B. Strajnar





















# Outline

- Cooperation with national provider
- Improved data set
- VarBC spin-up
- Impact experiments
- Conclusions
















## Cooperation with national provider

- Slovenian data (18 stations) not available through E-GVAP.
- ZTD solution computed by Geodetic Institute of Slovenia for around ~30 stations provided to ARSO (last few years)
- provided to ARSO in real time
- using Pivot software and provided to ARSO (last few years)





🧖 ZAMG
















# Old data set (Pivot)

- Earlier DA evaluations: clearly detrimental impact on humidity stations) not available through E-GVAP.
- ZTD solution computed by Geodetic Institute of Slovenia using Pivot software and provided to ARSO (last few years)




















# New data set (Bernese)

- GIS used Bernese geodetic software, I month of data provided (September 2019) for evaluation.
- GIS solution now within the spread of E-GVAP.



Evolution of ZTD estimate for Ljubljana (GSRI).














# New data set (Bernese)

- GIS used Bernese geodetic software, I month of data provided (September 2019) for evaluation.
- GIS solution now within the spread of E-GVAP.



Evolution of ZTD estimate for Ljubljana (GSRI).



ZAMG














# VarBC spin-up

- REF:A one-month assimilation cycle with ALADIN/SI(4.4 km)
- EXP: same setup using SIGNAL network, cold start of VarBC, using predictors 0 (constant) 1 (1000-300 hPa thickness), 4 (total column water) – default in cy43t2
- Non-negligible bias observed in REF (passive assimilation)


















# VarBC spin-up

**Bias correction evolution** 



Institute


OMSZ

DHMZ





### OMG in EXP: active assimilation



Mean: OMG 0,005 m (3-times less than in REF) Std.: 0.012 m (remains the same)



Czech Hydror

Institute















#### Recycled VarBC

# EXP2: same as EXP but recycled VarBC (warmed up by EXP)



STD of background departures



Mean: OMG 0,003 m (5-times less than in REF) Std.: 0.011 m (slightly drops)
















ARSO METEC Slovenia



#### Evolution of predictors – EXP1




#### Impact on forecast – upper air

- 2 production runs per day, 24h forecasts
- On average over domain, scores are neutral (small dataset)
- Over Slovenia, reduction of bias (T, RH) when using 3 closest radiosonde stations

No cases

No cases




Ра

<sup>3</sup> stations Selection: SIsurroundingsTEMP Temperature Period: 20190901-20191001 Statistics at 00 UTC Used {00,12} + 12 24

<sup>3</sup> stations Selection: SlsurroundingsTEMP Specific humidity Period: 20190901-20191001 Statistics at 00 UTC Used {00,12} + 12 24

#### Impact on forecast surface

- Clear degradation of specific humidity and also temperature
- More moisture, more clouds



100

25

20

-1.5


0

5

10


Forecast length

15





# Impact on forecast - precipitation

To much precipitation at 3 and 6 h range
EXP2 converges towards REF.




### Heavy precipitation case (+6h forecast)











SURFPREC.EAU.CON 2019/09/23 12:00 +6

#### EXP - REF





#### Conclusions

- Revised data set very close to E-GVAP, similar behavior expected
- ZTD have large impact on humidity and precipitation (should obs. error be tuned?)
- Bias correction VarBC seems to need longer period (3-5 months?)
- Should the static BC also be used and then variational?
- Cycling of VarBC (3h or 24h); how to technically handle?
- Continuation depends on agreement with GIS & E-GVAP